

# HONG KONG RNA CLUB

## Seminar



香港城市大學  
City University of Hong Kong

19 Jun 2018 (Tue) / 5:00-6:00pm

Room 2-130, To Yuen Building (桃源樓)

City University of Hong Kong

### Guest Speaker:

Prof. Ai Kotani  
School of Medicine  
Tokai University



*Role of secretory small RNAs  
as a pro-inflammatory mediator  
in the development of  
EBV-associated lymphoma*

### Sponsors:



**EXIQON**  
Seek Find Verify

**LEXOGEN**  
Enabling complete transcriptome sequencing

**BIO-RAD**

**ThermoFisher**  
SCIENTIFIC

**eppendorf**



**Refreshment provided!**

Sign-up: [kitkwok.com/hk-rna-club](http://kitkwok.com/hk-rna-club)

Contact: [hkrnaclub@gmail.com](mailto:hkrnaclub@gmail.com)

# Role of secretary small RNAs as a pro-inflammatory mediator in the development of EBV-associated lymphoma

**Professor Ai Kotani**

Department of Hematology and Oncology

Tokai University School of Medicine

Tokai University Institute of Innovative Science and Technology



## Abstract

Epstein-Barr virus (EBV) causes various diseases in the elderly including B-cell lymphoma such as Hodgkin's lymphoma (HL) and diffuse large B-cell lymphoma (DLBCL). Here, we show that EBV acts *in trans* on non-infected macrophages in the tumor through exosome secretion and augments the development of lymphomas. In a humanized mouse model, the different formation of lymphoproliferative disease (LPD) between two EBV strains (Akata and B95-8) was evident. Furthermore, injection of Akata derived exosomes affected LPD severity possibly through the regulation of macrophage phenotype *in vivo*. Exosomes collected from Akata- lymphoblastoid cell lines (LCLs) reportedly contain EBV-derived non-coding RNAs such as BamHI fragment A rightward transcript (BART) miRNAs and EBV-encoded RNA (EBER). We focused on the exosome-mediated delivery of BART miRNAs. *In vitro*, BART miRNAs could induce the immune regulatory phenotype in macrophages characterized by the gene expressions of *interleukin-10*, *tumor necrosis factor-alpha*, and *arginase 1*, suggesting the immune regulatory role of BART miRNAs. The expression level of an EBV-encoded miRNA was strongly linked to the clinical outcomes in elderly diffuse large B-cell lymphoma patients. These results implicate BART miRNAs as one of the factors regulating the severity of lymphoproliferative disease and as a diagnostic marker for EBV+ B-cell lymphoma.